arXiv:math/0002170v1 [math.QA] 21 Feb 2000

SYMMETRIZER AND ANTISYMMETRIZER OF THE BIRMAN–WENZL–MURAKAMI ALGEBRAS

ISTVÁN HECKENBERGER AND AXEL SCHÜLER

Abstract. Explicit formulas for the symmetrizer and the antisymmetrizer of the Birman–Wenzl–Murakami algebras $\mathrm{BWM}(r,q)_n$ are given.

AMS subject classification: 20C15, 20F36, 16G10

1. Introduction

The Birman–Wenzl–Murakami algebra was first defined and independently studied by Birman and Wenzl [1] and Murakami [4]. The Iwahori–Hecke algebras of Type A and the Birman–Wenzl–Murakami algebras naturally arise as centralizer algebras of tensor product corepresentations of quantum groups of Type A and of Type B, C, and D, respectively [5, 6]. Irreducible characters and primitive idempotents of Hecke algebras have been intensively studied during the last 30 years. There are explicit constructions of analogues of the Young symmetrizer and Specht modules [2, 3]. However, in case of the Birman–Wenzl–Murakami algebras there are only a few papers in this direction [1, 3]. Young symmetrizers for the Birman–Wenzl–Murakami algebra are not known.

The aim of this Letter is to give recursive but explicit formulas for the most important minimal central idempotents of the Birman–Wenzl–Murakami algebras namely for the symmetrizer S_n and the antisymmetrizer A_n. For the symmetrizer, we establish the formulas

$$cS_n = S_{n-1}b^+_{n-1,1} = a^+_{n-1,1}S_{n-1} = S_{n-1}[1]b^+_{1,n-1} = a^+_{n-1,1}S_{n-1}[1],$$

where c is a normalization constant. Here $S_{n-1}[1]$ denotes the symmetrizer S_{n-1} shifted by one. The elements $a^+_{1,n-1}$ and $a^+_{n-1,1}$ are the substitutes for the well-known sums of shuffle permutations

$$1+s_1+s_1s_2+\cdots+s_1\cdots s_{n-1} \quad \text{and} \quad 1+s_{n-1}+s_{n-1}s_{n-2}+\cdots+s_{n-1}\cdots s_1$$

in the group algebra of the symmetric group. The elements $b^+_{n-1,1}$ and $b^+_{1,n-1}$ correspond to the sum of the inverse shuffle permutations. Taking the quotient of $\mathrm{BWM}(r,q)_n$ by

---

Supported by the Deutsche Forschungsgemeinschaft.

the relations $e_1 = \cdots = e_{n-1} = 0$ one gets the well-known formula for the symmetrizer of the Hecke algebra

$$S_n = q^{-\frac{n(n-1)}{2}}([\![n]\!]!)^{-1} \sum_w q^{\ell(w)} T_w,$$

where the sum is over all elements w of the symmetric group of n variables. Here T_w denotes the corresponding elements of the Hecke algebra and $\ell(w)$ is the length of w. Similar formulas for the antisymmetrizer A_n are also given.

2. The Birman–Wenzl–Murakami Algebra

Let $n \in \mathbb{N}$, $n \geq 2$, and $r, q \in \mathbb{C} \setminus \{0\}$. Further, we assume that $q^k \neq 1$ for any $k \in \mathbb{N}$ and $r \neq \pm q^k$ for any $k \in \mathbb{Z}$. We use the abbreviation $\hat{q}$ for the complex number $q - q^{-1}$. The symbol $[\![k]\!]$, $k \in \mathbb{Z}$, denotes the complex number $(q^k - q^{-k})/(q - q^{-1})$.

The Birman–Wenzl–Murakami algebra $\mathrm{BWM}(r,q)_n$ is the unital algebra over the complex numbers generated by the elements g_i, e_i, $i = 1, \ldots, n-1$, and relations

$$g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1}, \qquad g_i g_j = g_j g_i \quad \text{for } |i-j| > 1, \tag{1}$$

$$g_i^2 = 1 + \hat{q} g_i - r^{-1}\hat{q} e_i, \qquad g_i e_i = e_i g_i = r^{-1} e_i, \tag{2}$$

$$g_i g_{i+1} e_i = e_{i+1} e_i, \qquad g_{i+1} g_i e_{i+1} = e_i e_{i+1}. \tag{3}$$

For any $n \in \mathbb{N}$, $n \geq 2$, the algebra $\mathrm{BWM}(r,q)_n$ is naturally embedded into the algebras $\mathrm{BWM}(r,q)_{n+k}$, $k \geq 0$.

There is an algebra automorphism α_n of $\mathrm{BWM}(r,q)_n$ defined by $\alpha_n(g_i) = g_{n-i}$, $\alpha_n(e_i) = e_{n-i}$ for all $i = 1, \ldots, n-1$. There is an algebra antiautomorphism β_n of $\mathrm{BWM}(r,q)_n$ defined by $\beta_n(g_i) = g_i$, $\beta_n(e_i) = e_i$ for any $i = 1, \ldots, n-1$. There is an algebra isomorphism $\gamma_n : \mathrm{BWM}(r,q)_n \to \mathrm{BWM}(r,p)_n$, $p = -q^{-1}$, given by $\gamma_n(g_i) = g_i$, $\gamma_n(e_i) = e_i$.

Let $s : \mathrm{BWM}(r,q)_n \to \mathrm{BWM}(r,q)_{n+1}$ be the algebra homomorphism defined by $s(g_i) := g_{i+1}$, $s(e_i) = e_{i+1}$. If $b =: b[0] \in \mathrm{BWM}(r,q)_n$ and $s^k(b) \in \mathrm{BWM}(r,q)_n$ for some $k \in \mathbb{N}$, then we write $b[k]$ for the element $s^k(b)$.

3. The Symmetrizer

The symmetrizer S_n is the unique nonzero element of the algebra $\mathrm{BWM}(r,q)_n$ such that $S_n^2 = S_n$ and $S_n g_i = q S_n$ for any $i = 1, \ldots, n-1$ [3, Theorem 5.14]. It follows that S_n is central, in particular $g_i S_n = q S_n$ for all $i = 1, \ldots, n-1$. It is well known that

$$S_2 = \frac{1}{q[\![2]\!]}\left(1 + q g_1 + \frac{q\hat{q}}{1 - qr} e_1\right). \tag{4}$$

Now let us introduce the elements

$$d^+_{k,i} := e_{k-1}e_{k-2}\cdots e_i \sum_{j=0}^{i-1} q^j g_{i-1}\cdots g_{i-j}, \qquad 1 \le i < k, \tag{5}$$

$$b^+_{k,1} := \sum_{i=0}^{k} q^i g_k g_{k-1}\cdots g_{k+1-i} + \frac{\hat{q}}{1-q^{2k-1}r}\sum_{i=1}^{k} q^{2k-2i+1} d^+_{k+1,i}, \qquad 0 \le k < n. \tag{6}$$

More precisely, formula (5) reads as $d^+_{k,i} = e_{k-1}e_{k-2}\cdots e_i \sum_{j=0}^{i-1} q^j \prod_{l=1}^{j} g_{i-l}$.

Proposition 1. *For any $n \ge 3$ the formula $S_n = S_{n-1}b^+_{n-1,1}/(q^{n-1}[\![n]\!])$ holds.*

Lemma 2. *The expression $S_{n-1}d^+_{n,k}g_l$, $l < n$, equals*

$$\begin{aligned}
&qS_{n-1}d^+_{n,k} - qr^{-1}\hat{q}S_{n-1}e_{n-1}e_{n-2}\cdots e_l && \text{for } l < k,\\
&qS_{n-1}d^+_{n,k-1} + r^{-1}S_{n-1}e_{n-1}e_{n-2}\cdots e_l && \text{for } l = k > 1,\\
&r^{-1}S_{n-1}d^+_{n,k} && \text{for } l = k = 1,\\
&q^{-1}S_{n-1}d^+_{n,k+1} + \hat{q}S_{n-1}d^+_{n,k} - q^{2k-1}S_{n-1}e_{n-1}e_{n-2}\cdots e_l && \text{for } l = k+1,\\
&qS_{n-1}d^+_{n,k} && \text{for } l \ge k+2.
\end{aligned}$$

Proof of the Proposition. Let $S'_n := S_{n-1}b^+_{n-1,1}/(q^{n-1}[\![n]\!])$. We prove that $S'_n g_i = qS'_n$ for any $i = 1, 2, \ldots, n-1$. Then we conclude that $S'_n e_i = q^{-1}S'_n g_i e_i = q^{-1}r^{-1}S'_n e_i$ and hence $(q^{-1}r^{-1} - 1)S'_n e_i = 0$. Since $r \ne q^{-1}$, we obtain $S'_n e_i = 0$. Now (4) and the assertion of the Proposition for $k < n$ imply that

$${S'_n}^2 = S'_n S_2 b^+_{2,1} b^+_{3,1}\cdots b^+_{n-1,1}/(q^2[\![3]\!]q^3[\![4]\!]\cdots q^{n-1}[\![n]\!]) = S'_n. \tag{7}$$

Hence, S'_n is the symmetrizer of the algebra $\mathrm{BWM}(r,q)_n$.

Let $\tilde{S}_n = S_{n-1}b^+_{n-1,1}$. We prove that $\tilde{S}_n g_i = q\tilde{S}_n$ for any $i = 1, 2, \ldots, n-1$. For this we use the relations of the algebra $\mathrm{BWM}(r,q)_n$ and Lemma 2. Observe that in Lemma 2 only S_{n-1} occurs. Therefore, we can use that $S_{n-1}g_i = qS_{n-1}$ for all $i = 1, \ldots, n-2$. Let now $i \ge 2$. Then

$$\begin{aligned}
\tilde{S}_n g_i =& S_{n-1}\left(\sum_{j=0}^{n-1} q^j g_{n-1}g_{n-2}\cdots g_{n-j} + \frac{\hat{q}}{1-q^{2n-3}r}\sum_{j=1}^{n-1} q^{2n-2j-1}d^+_{n,j}\right) g_i\\
=& S_{n-1}\Bigg(q\sum_{j=0}^{n-i-2} q^j g_{n-1}\cdots g_{n-j} + q^{n-i-1}g_{n-1}\cdots g_{i+1}g_i +\\
&+ q^{n-i}g_{n-1}\cdots g_{i+1}(1 + \hat{q}g_i - r^{-1}\hat{q}e_i) + q\sum_{j=n-i+1}^{n-1} q^j g_{n-1}\cdots g_{n-j} +
\end{aligned}$$

$$
\begin{aligned}
&+\frac{\hat{q}}{1-q^{2n-3}r}\Bigg(\sum_{j=1}^{i-2} q^{2n-2j-1} q d^+_{n,j} + q^{2n-2i+1}(q^{-1} d^+_{n,i} + \hat{q} d^+_{n,i-1} - \\
&- q^{2i-3} e_{n-1}\cdots e_i) + q^{2n-2i-1}(q d^+_{n,i-1} + r^{-1} e_{n-1}\cdots e_i) + \\
&+ \sum_{j=i+1}^{n-1} q^{2n-2j-1}(q d^+_{n,j} - q r^{-1}\hat{q} e_{n-1}\cdots e_i)\Bigg)\Bigg) \\
=& S_{n-1}\Bigg(q\sum_{j=0}^{n-1} q^j g_{n-1}\cdots g_{n-j} - q^{n-i} q^{-n+i+1} r^{-1}\hat{q} e_{n-1}\cdots e_{i+1} e_i + \\
&+\frac{\hat{q}}{1-q^{2n-3}r}\Bigg(\sum_{j=1}^{n-1} q^{2n-2j} d^+_{n,j} + (-q^{2n-2} + q^{2n-2i-1} r^{-1} - \\
&- r^{-1}\hat{q} q^{n-i} [\![n-1-i]\!]) e_{n-1}\cdots e_i\Bigg)\Bigg) \\
=& q S_{n-1} b^+_{n-1,1} + \hat{q} S_{n-1}(-q r^{-1} + (-q^{2n-2} + r^{-1} q)/(1-q^{2n-3} r)) e_{n-1}\cdots e_i \\
=& q\tilde{S}_n.
\end{aligned}
$$

The case $i=1$ can be proven similarly. □

Proof of the Lemma. Let $l<k$. Using the relations (1)–(3) we obtain

$$
\begin{aligned}
S_{n-1} d^+_{n,k} g_l =& S_{n-1} e_{n-1}\cdots e_k \sum_{i=0}^{k-1} q^i g_{k-1}\cdots g_{k-i} g_l \\
=& S_{n-1} e_{n-1}\cdots e_k \Bigg(q\sum_{i=0}^{k-l-2} q^i g_{k-1}\cdots g_{k-i} + q^{k-l-1} g_{k-1}\cdots g_{l+1} g_l + \\
&+ q^{k-l} g_{k-1}\cdots g_{l+1}(1 + \hat{q} g_l - r^{-1}\hat{q} e_l) + q\sum_{i=k-l+1}^{k-1} q^i g_{k-1}\cdots g_{k-i}\Bigg) \\
=& S_{n-1} e_{n-1}\cdots e_k \left(q\sum_{i=0}^{k-1} q^i g_{k-1}\cdots g_{k-i} - q^{k-l} r^{-1}\hat{q} q^{-k+l+1} e_{k-1}\cdots e_l\right) \\
=& q S_{n-1} d^+_{n,k} - q r^{-1}\hat{q} S_{n-1} e_{n-1} e_{n-2}\cdots e_l.
\end{aligned}
$$

The other cases can be shown similarly. □

If we apply the automophism $\varphi := \alpha_n$ (the antiautomorphisms $\varphi := \beta_n$ and $\varphi := \alpha_n \circ \beta_n$, respectively,) to the symmetrizer S_n, we reobtain S_n. Indeed, the image of S_n is an idempotent, $\varphi(S_n)\varphi(S_n) = \varphi(S_n^2) = \varphi(S_n)$. Further,

$$
\varphi(S_n) g_i = \varphi(S_n g_{n-i}) = q\varphi(S_n) \qquad \text{for all } i = 1,\dots,n-1 \tag{8}
$$

($\varphi(S_n)g_i = \varphi(\varphi(g_i)S_n) = \varphi(qS_n)$ for all $i = 1, \dots, n-1$). Thus we obtain the following formulas for the symmetrizer S_n:

$$d'^{+}_{k,i} := \alpha_k(d^{+}_{k,i}) = e_1 e_2 \cdots e_{k-i} \sum_{j=0}^{i-1} q^j g_{k+1-i} \cdots g_{k-i+j}, \tag{9}$$

$$b^{+}_{1,k} := \alpha_{k+1}(b^{+}_{k,1}) = \sum_{i=0}^{k} q^i g_1 g_2 \cdots g_i + \frac{\hat{q}}{1 - q^{2k-1} r} \sum_{i=1}^{k} q^{2k-2i+1} d'^{+}_{k+1,i}, \tag{10}$$

$$\bar{d}^{+}_{k,i} := \beta_n(d^{+}_{k,i}) = \sum_{j=0}^{i-1} q^j g_{i-j} \cdots g_{i-1} e_i e_{i+1} \cdots e_{k-1}, \tag{11}$$

$$a^{+}_{k,1} := \beta_n(b^{+}_{k,1}) = \sum_{i=0}^{k} q^i g_{k+1-i} \cdots g_{k-1} g_k + \frac{\hat{q}}{1 - q^{2k-1} r} \sum_{i=1}^{k} q^{2k-2i+1} \bar{d}^{+}_{k+1,i}, \tag{12}$$

$$\bar{d}'^{+}_{k,i} := \beta_n(d'^{+}_{k,i}) = \sum_{j=0}^{i-1} q^j g_{k-i+j} \cdots g_{k-i+1} e_{k-i} \cdots e_2 e_1, \tag{13}$$

$$a^{+}_{1,k} := \beta_n(b^{+}_{1,k}) = \sum_{i=0}^{k} q^i g_i \cdots g_2 g_1 + \frac{\hat{q}}{1 - q^{2k-1} r} \sum_{i=1}^{k} q^{2k-2i+1} \bar{d}'^{+}_{k+1,i}. \tag{14}$$

$$\begin{aligned} S_n &= S_{n-1} b^{+}_{n-1,1} / (q^{n-1} [\![n]\!]) = S_{n-1}[1] b^{+}_{1,n-1} / (q^{n-1} [\![n]\!]) \\ &= a^{+}_{n-1,1} S_{n-1} / (q^{n-1} [\![n]\!]) = a^{+}_{1,n-1} S_{n-1}[1] / (q^{n-1} [\![n]\!]). \end{aligned} \tag{15}$$

4. THE ANTISYMMETRIZER

It is well known that the antisymmetrizer A_n is the unique nonzero idempotent in the algebra $\mathrm{BWM}(r,q)_n$ such that $A_n g_i = -q^{-1} g_i$ for all $i = 1, \dots, n-1$. Let us examine the image A'_n of the symmetrizer S_n under the isomorphism $\gamma_n : \mathrm{BWM}(r,p)_n \to \mathrm{BWM}(r,q)_n$, $p = -q^{-1}$ [6, Proposition 3.2 (c)]. Obviously, ${A'_n}^2 = \gamma_n(S_n)^2 = \gamma_n(S_n^2) = \gamma_n(S_n) = A'_n$, hence A'_n is an idempotent. Moreover,

$$A'_n g_i = \gamma_n(S_n)\gamma_n(g_i) = \gamma_n(S_n g_i) = p\gamma_n(S_n) = -q^{-1} A'_n \tag{16}$$

for all $i = 1, \dots, n-1$. The explicit formulas for A'_n show that A'_n is nonzero. Hence, A'_n is the antisymmetrizer of the algebra $\mathrm{BWM}(r,q)_n$. We obtain the following formulas:

$$d^{-}_{k,i} := \gamma_n(d^{+}_{k,i}) = e_{k-1} e_{k-2} \cdots e_i \sum_{j=0}^{i-1} (-q)^{-j} g_{i-1} \cdots g_{i-j}, \tag{17}$$

$$b^{-}_{k,1} := \gamma_n(b^{+}_{k,1}) = \sum_{i=0}^{k} (-q)^{-i} g_k g_{k-1} \cdots g_{k+1-i} - \frac{\hat{q}}{1 + q^{-2k+1} r} \sum_{i=1}^{k} q^{2i-2k-1} d^{-}_{k+1,i}, \tag{18}$$

$$d'^{-}_{k,i} := \gamma_n(d'^{+}_{k,i}) = e_1 e_2 \cdots e_{k-i} \sum_{j=0}^{i-1} (-q)^{-j} g_{k+1-i} \cdots g_{k-i+j}, \tag{19}$$

$$b^{-}_{1,k} := \gamma_n(b^{+}_{1,k}) = \sum_{i=0}^{k} (-q)^{-i} g_1 g_2 \cdots g_i - \frac{\hat{q}}{1+q^{-2k+1}r} \sum_{i=1}^{k} q^{2i-2k-1} d'^{-}_{k+1,i}, \tag{20}$$

$$\bar{d}^{-}_{k,i} := \gamma_n(\bar{d}^{+}_{k,i}) = \sum_{j=0}^{i-1} (-q)^{-j} g_{i-j} \cdots g_{i-1} e_i e_{i+1} \cdots e_{k-1}, \tag{21}$$

$$a^{-}_{k,1} := \gamma_n(a^{+}_{k,1}) = \sum_{i=0}^{k} (-q)^{-i} g_{k+1-i} \cdots g_{k-1} g_k - \frac{\hat{q}}{1+q^{-2k+1}r} \sum_{i=1}^{k} q^{2i-2k-1} \bar{d}^{-}_{k+1,i}, \tag{22}$$

$$\bar{d}'^{-}_{k,i} := \gamma_n(\bar{d}'^{+}_{k,i}) = \sum_{j=0}^{i-1} (-q)^{-j} g_{k-i+j} \cdots g_{k-i+1} e_{k-i} \cdots e_2 e_1, \tag{23}$$

$$a^{-}_{1,k} := \gamma_n(a^{+}_{1,k}) = \sum_{i=0}^{k} (-q)^{-i} g_i \cdots g_2 g_1 - \frac{\hat{q}}{1+q^{-2k+1}r} \sum_{i=1}^{k} q^{2i-2k-1} \bar{d}'^{-}_{k+1,i}. \tag{24}$$

For the antisymmetrizer $A_n = \gamma_n(S_n)$ the following equations hold:

$$A_n = q^{n-1} A_{n-1} b^{-}_{n-1,1} / \llbracket n \rrbracket = q^{n-1} A_{n-1}[1] b^{-}_{1,n-1} / \llbracket n \rrbracket \tag{25}$$

$$= q^{n-1} a^{-}_{n-1,1} A_{n-1} / \llbracket n \rrbracket = q^{n-1} a^{-}_{1,n-1} A_{n-1}[1] / \llbracket n \rrbracket. \tag{26}$$

REFERENCES

1. J.S. Birman and H. Wenzl, *Braids, link polynomials and a new algebra*, Trans. Amer. Math. Soc. **313** (1989), no. 1, 249–273.
2. A. Gyoja, *A q-analogue of Young symmetrizer*, Osaka J. Math. **23** (1986), 841–852.
3. R. Leduc and A. Ram, *A ribbon Hopf algebra approach to the irreducible representations of centralizer algebras: The Brauer, Birman-Wenzl, and type A Iwahori-Hecke algebras.*, Adv. Math. **125** (1997), no. 1, 1–94.
4. J. Murakami, *The representations of the q-analogue of Brauer's centralizer algebras and the Kauffman polynomial of links*, Publ. Res. Inst. Math. Sci. **26** (1990), no. 6, 935–945.
5. H. Wenzl, *Hecke algebras of type A_n and subfactors*, Invent. Math. **92** (1988), no. 2, 349–383.
6. ———, *Quantum groups and subfactors of type B, C, and D*, Commun. Math. Phys. **133** (1990), no. 2, 383–432.

UNIVERSITÄT LEIPZIG, MATHEMATISCHES INSTITUT, AUGUSTUSPLATZ 10, D-04109 LEIPZIG, GERMANY

E-mail address: `{heckenbe,schueler}@mathematik.uni-leipzig.de`